

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
30 June 2005 (30.06.2005)


PCT

(10) International Publication Number
WO 2005/059167 A1

- (51) International Patent Classification⁷: C12Q 1/44, (72) Inventors; and
G01N 33/68, C12Q 1/68 (75) Inventors/Applicants (for US only): PALLAORO,
Michele [IT/IT]; IRBM, Via Pontina KM30.600, I-00040
(21) International Application Number: PCT/EP2004/014159 Pomezia (IT). GALLINARI, Paola [IT/IT]; IRBM, Via
Pontina KM30.600, I-00040 Pomezia (IT). ALTAMURA,
Sergio [IT/IT]; IRBM, Via Pontina KM30.600, I-00040
(22) International Filing Date: Pomezia (IT). STEINKUHLER, Christian [DE/IT];
IRBM, Via Pontina KM30.600, I-00040 Pomezia (IT).
10 December 2004 (10.12.2004)
(25) Filing Language: English (74) Agent: MAN, Jocelyn; Merck & Co., Inc., European
Patent Department, Terlings Park, Eastwick Road, Harlow
(26) Publication Language: English Essex CM20 2QR (GB).
(30) Priority Data: 60/530,540 18 December 2003 (18.12.2003) US (81) Designated States (unless otherwise indicated, for every
kind of national protection available): AE, AG, AL, AM,
AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN,
CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI,
GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE,
KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD,
MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG,
(71) Applicant (for all designated States except US): ISTI-
TUTO DI RICERCHE DI BIOLOGIA MOLECO-
LARE P ANGELETTI SPA [IT/IT]; Via Pontina KM
30.600, I-00040 Pomezia (IT).

[Continued on next page]

(54) Title: METHOD FOR IDENTIFYING HISTONE DEACETYLASE INHIBITORS



(57) Abstract: A method for identifying analytes which are histone deacetylase (HDAC) inhibitors is described. The method provides cells comprising a reporter gene operably linked to a transcription regulatory sequence, which includes nucleotide sequences responsive to a known HDAC inhibitor or subtype, stably integrated into the genome of the cells. A culture of the cells is incubated in medium which contains an analyte and the culture monitored for expression of the reporter gene. Analytes which have HDAC inhibitor activity induce expression of the reporter gene. In particular embodiments, the transcription regulatory region is a sequence of the p21WAF1/CIP1 transcription regulatory sequence which is responsive to a known HDAC inhibitor but not responsive to p53.

WO 2005/059167 A1



PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(84) Designated States (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, MC, NL, PL, PT, RO,

Published:

— with international search report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.